

Are you ready for depression symptom relief that's

FAST & LASTS

Some people saw depression symptom relief in as early as 1 week vs placebo. On average, significant symptom relief was seen at 6 weeks.*

If you're experiencing depression symptoms or treatment side effects, it's important to talk to your healthcare provider to help find a treatment option for you. This quick guide can help you learn more about AUVELITY.

*Measured against placebo on a depression rating scale during a 6-week study of 327 adults with MDD.
MDD=major depressive disorder.

WHAT IS AUVELITY (aw-VEHL-ah-tee)?

It is a prescription oral medicine used to treat adults with major depressive disorder (MDD). It is not known if AUVELITY is safe and effective for use in children.

AUVELITY is not approved for uses other than the treatment of MDD. The ingredients in AUVELITY, bupropion and dextromethorphan, are the same ingredients found in some other medicines approved for other uses.

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT AUVELITY?

AUVELITY and other antidepressant medicines may increase suicidal thoughts and actions in some children, adolescents, and young adults, especially within the first few months of treatment or when the dose is changed. AUVELITY is not for use in children.

You should pay close attention to any new or sudden changes in mood, behavior, thoughts, or feelings or if you develop suicidal thoughts or actions. This is very important when starting or changing the dose of an antidepressant medicine.

Please see full Important Safety Information throughout, full [Prescribing Information](#), including **Boxed Warning** for suicidal thoughts and behaviors, and [Medication Guide](#).

What if you could get your GO back?

 **Auvelity**[®]
(dextromethorphan HBr and bupropion HCl)
extended-release tablets 45mg/105mg

Talk to your healthcare provider about all the depression symptoms or treatment side effects you're experiencing (if you're taking medication right now).

I've been:

- Sad or down
- Not interested in things I used to like
- Low on energy or more tired
- Feeling worthless or guilty
- Having appetite changes (I've lost or gained weight, and it's not linked to dieting)
- Sleeping too much or having a hard time sleeping
- Having a hard time concentrating, thinking, or making decisions
- Moving/talking more slowly, or restless and fidgety (and others have noticed)
- Thinking about death or suicide
- Other

If I've experienced side effects on my current medication, they've been:

- Nausea
- Vomiting
- Diarrhea
- Sexual problems
- Weight gain
- Feeling sleepy and/or trouble sleeping
- Other

Note: Don't stop taking an antidepressant without talking to your healthcare provider first.

If you're thinking about suicide, call the confidential National Suicide Prevention Lifeline 24/7, at 1-800-273-TALK (8255). If you're having a medical emergency, call 911.

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT AUVELITY? (CONT'D)

Call your healthcare provider (HCP) or get emergency help right away if you or your loved one have any of the following symptoms, especially if they are new, worse, or worry you:

- suicidal thoughts or actions
- new or worsening depression or anxiety
- agitation or restlessness
- trouble sleeping (insomnia)
- acting aggressive, being angry, or violent
- an extreme increase in activity and talking (mania)
- panic attacks
- new or worsening irritability
- acting on dangerous impulses
- other unusual changes in behavior or mood

Do not take AUVELITY if you:

- have or had a seizure disorder.
- have or had an eating disorder like anorexia or bulimia.
- have recently and suddenly stopped drinking alcohol or use medicines called benzodiazepines, barbiturates, or anti-seizure medicines, and you have recently suddenly stopped taking them.
- are taking a monoamine oxidase inhibitor (MAOI), have stopped taking an MAOI in the last 14 days, or are being treated with the antibiotic linezolid or intravenous methylene blue. Ask your HCP or pharmacist if you are unsure whether you take an MAOI. Do not start taking an MAOI until you have stopped taking AUVELITY for at least 14 days.
- are allergic to dextromethorphan, bupropion, or any other ingredients in AUVELITY.

AUVELITY may cause serious side effects. Ask your HCP how to recognize the serious side effects below and what to do if you think you have one:

Seizures. There is a risk of seizures during treatment with AUVELITY. The risk is higher if you take higher doses of AUVELITY, have certain medical problems, or take AUVELITY with certain other medicines. **Do not** take AUVELITY with other medicines unless your healthcare provider tells you to.

If you have a seizure during treatment with AUVELITY, stop taking AUVELITY and call your HCP right away. **Do not** take AUVELITY again if you have a seizure.

Increases in blood pressure (hypertension). Some people may get high blood pressure during treatment with AUVELITY. Your HCP should check your blood pressure before you start taking and during treatment with AUVELITY.

Manic episodes. Manic episodes may happen in people with bipolar disorder who take AUVELITY. Symptoms may include:

- greatly increased energy
- racing thoughts
- unusually grand ideas
- talking more or faster than usual
- severe trouble sleeping
- reckless behavior
- excessive happiness or irritability

Unusual thoughts or behaviors. One of the ingredients in AUVELITY (bupropion) can cause unusual thoughts or behaviors, including delusions (believing you are someone else), hallucinations (seeing or hearing things that are not there), paranoia (feeling that people are against you), or feeling confused. If this happens to you, call your HCP.

Please see full Important Safety Information throughout, full [Prescribing Information](#), including **Boxed Warning** for suicidal thoughts and behaviors, and [Medication Guide](#).

Ready to learn more about AUVELITY?

Auvelity[®]
(dextromethorphan HBr and bupropion HCl)
extended-release tablets 45mg/105mg

Here are some good questions to ask your healthcare provider:



- What makes AUVELITY different from other antidepressants?
- How early can AUVELITY start working?
- What are the potential side effects?
- How do I take AUVELITY? How can I tell it's working?
- Can I take AUVELITY with my other medications?

Notes:

(other questions you have or things you want to mention)

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT AUVELITY? (CONT'D)

Eye problems (angle-closure glaucoma). AUVELITY may cause a type of eye problem called angle-closure glaucoma in people with certain other eye conditions. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are. Call your HCP if you have eye pain, changes in your vision, or swelling or redness in or around the eye.

Dizziness. AUVELITY may cause dizziness which may increase your risk for falls.

Serotonin syndrome. A potentially life-threatening problem called serotonin syndrome can happen when you take AUVELITY with certain other medicines. **Call your HCP or go to the nearest hospital emergency room right away** if you have any of the following signs and symptoms:

- agitation
- hallucinations
- confusion
- coma
- fast heartbeat
- blood pressure changes
- dizziness
- sweating
- flushing
- high body temperature (hyperthermia)
- shaking (tremors), stiff muscles, or muscle twitching
- loss of coordination
- seizures
- nausea, vomiting, diarrhea

COMMON SIDE EFFECTS

The most common side effects of AUVELITY include dizziness, headache, diarrhea, feeling sleepy, dry mouth, sexual function problems, and excessive sweating.

These are not all the possible side effects of AUVELITY. Tell your doctor if you have any side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see full Important Safety Information throughout, full [Prescribing Information](#), including **Boxed Warning** for suicidal thoughts and behaviors, and [Medication Guide](#).


BEFORE USING

- Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.
- It is important to tell your HCP if you are taking:
 - other medicines containing bupropion or dextromethorphan
 - medicines to treat depression, anxiety, psychotic or thought disorders, including selective serotonin reuptake inhibitors (SSRIs) and tricyclic antidepressants
 - theophylline
 - corticosteroids
 - oral diabetes medicines or use insulin to control your blood sugar
 - medicines to control appetite (anorectic)
 - nicotine medicines to help you stop smoking
 - street (illicit) drugs
 - benzodiazepines, sedative-hypnotic (sleep medicines), or opiates
- If you are unsure if you take any of these medicines, ask your HCP. They can tell you if it is safe to take AUVELITY with your other medicines.
- Tell your HCP if you are pregnant or plan to become pregnant. AUVELITY may harm your unborn baby if you take it during pregnancy. AUVELITY is not recommended during pregnancy. Your HCP will prescribe another treatment for females who plan to become pregnant.
- One of the ingredients in AUVELITY passes into your breast milk. Do not breastfeed during treatment with AUVELITY and for 5 days after the final dose.

You are not alone

Real patients have found depression symptom relief with AUVELITY.

// I got my game face back. //



Danielle. Real AUVELITY patient since 2023.



Scan or visit [AUVELITY.com](https://www.auvelity.com) to learn more about savings and support options.

BEFORE USING (CONT'D)

Tell your HCP about all your medical conditions, including if you:

- have problems with your liver or kidneys.
- have diabetes, heart disease, or high blood pressure.
- have a history of seizure, stroke, eating disorder, head injury, or have a tumor in your brain or spinal cord.
- have a history of alcohol or drug abuse.
- have a history of seizure, eating disorder, or abuse alcohol or drugs.
- have low blood sugar, low blood sodium levels, or a history of falls.
- you take certain other medicines that could interact with AUVELITY.
- have or had a condition known as bipolar disorder, a family history of bipolar disorder, suicide, or depression.
- have high pressure in the eye (glaucoma).

Review the list below with your HCP. AUVELITY may not be right for you if:

- you drink a lot of alcohol.
- you abuse prescription or street drugs.
- you are pregnant or plan to become pregnant.
- you are breastfeeding or plan to breastfeed.

HOW TO TAKE

- AUVELITY is available by prescription only.
- Take AUVELITY exactly as instructed by your HCP.

Please see full Important Safety Information throughout, full [Prescribing Information](#), including **Boxed Warning** for suicidal thoughts and behaviors, and [Medication Guide](#).

- Take AUVELITY 1 time a day for 3 days, then increase your dose to 2 times a day (taken at least 8 hours apart). Do not take more than 2 AUVELITY tablets in 24 hours.
- If you miss a dose, do not take an extra dose. Wait and take your next dose at the regular time. **Do not** take more than 1 dose of AUVELITY at a time.
- Do not change your dose or stop taking AUVELITY without talking to your HCP.
- Swallow AUVELITY tablets whole. Do not crush, chew, or divide the tablets.
- Do not give AUVELITY to other people.
- If you take too much AUVELITY call your poison control center at 1-800-222-1222 or go to the nearest hospital emergency room right away.

LEARN MORE

For more information about AUVELITY, call 866-496-2976 or visit [AUVELITY.com](https://www.auvelity.com).

This summary provides basic information about AUVELITY but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your doctor. Be sure to talk to your doctor or other HCP about AUVELITY and how to take it. Your HCP is the best person to help you decide if AUVELITY is right for you.

AUV CON BS 08/2024

axsome

AUVELITY, AXSOME, and their logos are trademarks or registered trademarks of Axsome Therapeutics, Inc. or its affiliates. Other trademarks are property of their respective owners. © 2025 Axsome Therapeutics, Inc. All rights reserved. PP-AUV-US-2100028 04/2025

 **Auvelity**[®]
(dextromethorphan HBr and bupropion HCl)
extended-release tablets 45mg/105mg